

COMPARATIVE STUDY OF THE ANTIBACTERIAL, ANTI-INFLAMMATORY, AND ANTIOXIDANT EFFECTS OF HONEY, PROPOLIS, AND ROYAL JELLY PRODUCED BY THE *APIS MELLIFERA* BEE

Mabrouka Bouacha^{1,2}  Ines Boudiar¹ 

¹Laboratory of Microbiology and Molecular Biology, Sciences Faculty, Badji Mokhtar University, Annaba, Algeria ²Laboratory of Biochemistry and Environmental Toxicology, Sciences Faculty, Badji Mokhtar University, Annaba, Algeria

Honey, propolis, and royal jelly are natural products of bees that have been used for medicinal purposes for centuries. The aim of this study is to investigate and compare the therapeutic effects of two samples of honey, two of propolis, and two of royal jelly. All hive products were produced by the *Apis mellifera* bee and collected from two regions of Algeria. The study evaluated the antibacterial activity of hive products using agar well-diffusion and microbroth dilution assays. The agar well-diffusion assay involved inoculating agar plates with ten different bacterial strains, while the micro-broth dilution assay involved serial dilutions of hive products and bacterial cultures. The antioxidant potential was assessed using scavenging assays of 2,2-diphenyl-1-picrylhydrazyl (DPPH) and ferric-reducing antioxidant power (FRAP) assays, which measured the product's ability to donate electrons and neutralize DPPH free radicals. The albumin denaturation method evaluated the anti-inflammatory properties of the hive products by measuring the extent of inhibition of protein denaturation under heat stress. The results demonstrated significant differences in the antibacterial effects of the tested products; honey samples showed greater antibacterial activity than propolis and royal jelly samples. Gram-positive bacteria were more susceptible to the antibacterial effects than Gram-negative bacteria. Regarding antioxidant and anti-inflammatory properties, honey samples were more effective than propolis and royal jelly samples. Propolis sample 1 had the highest concentration of polyphenols and flavonoids. All hive products exhibited potent antibacterial, antioxidant, and anti-inflammatory effects, making them a promising natural supplement for combating various health issues associated with bacterial infections, oxidative stress, and inflammation.

Keywords: *Apis mellifera*, honey, propolis, royal jelly, therapeutic properties

Submitted: April 7, 2023 **Accepted:** November 20, 2025

Published online: March 15, 2026

Copyright: © 2026, Author(s). This is an open-access article published under the terms of the Creative Commons Attribution 4.0 International License. (<http://creativecommons.org/licenses/by/4.0/>).

Correspondence to:

Mabrouka Bouacha
Laboratory of Microbiology and Molecular Biology
Sciences Faculty, Badji Mokhtar University
Annaba, Algeria
E-mail: bouacha.mabrouka@yahoo.fr

INTRODUCTION

The *Apis mellifera* bee is a social insect belonging to the genus *Apis* and the family *Apidae*. It is of considerable economic and cultural importance for human society as its role in crop pollination is essential for global food production (1). In addition, the *Apis mellifera* bee provides natural products such as propolis, pollen, venom, royal jelly, and honey with many potential healthy qualities, including antimicrobial (2,3), antimutagenic (4), antioxidant (5), and antitumoral effects (6). Commercialization and export of *Apis mellifera*'s products are a source of income for several countries worldwide, especially for honey production (1,7). Moreover, due to their composition of several bioactive substances, natural bee products are increasingly used in research to develop new treatment alternatives for several diseases (8).

Several properties of honey contribute to its antimicrobial effects. It has a pH between 3.2 and 4.5, which inhibits the growth of most pathogenic microorganisms (9,10). It contains a very high concentration of sugar (glucose and fructose), which dehydrates bacteria and other microorganisms. It also contains glucose oxidase, an enzyme that produces hydrogen peroxide (H_2O_2), which is responsible for the inhibition of bacterial growth. Other factors contribute to the antibacterial effect of honey, such as lysozymes, flavonoids, polyphenols, and other aromatic substances derived from foraged plants (11). In addition, honey has antioxidant activity, preventing oxidation by neutralizing free radicals in the body, which can damage cells, including DNA being also responsible for cell aging. The antioxidant properties of honey are attributed to many biological substances, such as phenolic acids, flavonoids, vitamins, enzymes, and minerals (12). Moreover, honey could also have anti-inflammatory and antioxidant properties, which are associated with each other through multiple cellular signaling pathways (13). Honey has been found to inhibit inflammation, oxidative stress, and gene expression pathways. Inflammation is a crucial part of the body's immune response, but chronic inflammation can lead to diseases like arthritis and cardiovascular disorders. Honey's polyphenolic and flavonoid content can modulate these processes by acting as agonists or inhibitors of key pro-inflammatory cytokines. Honey's antioxidants help counteract oxidative stress by scavenging free radicals and enhancing the body's antioxidant defenses. By addressing both inflammation and oxidative stress, honey not only addresses individual issues but also disrupts the cycle that perpetuates their coexistence, contributing to its

therapeutic potential in preventing and managing chronic inflammatory and oxidative stress-related diseases (5,6,13).

Royal jelly is the most elaborate substance of the hive. It is intended for the feeding of queens. It is a creamy, whitish-to-yellowish secretion, composed of water (about 66%), carbohydrates (about 14.5%), fatty acids (about 4.5%), including essential fatty acids, and proteins (about 13%), as well as essential amino acids. The antibacterial activity of royal jelly is mainly due to its acidic pH, trans-10-hydroxy-2-decenoic acid, and proteins with glucose-oxidase activity, such as major royal jelly proteins, defensin, royalisin, jelleins, and apisimin (14,15).

Propolis, a resinous substance collected by bees from various botanical sources, is used as a building material to fortify hives and protect them from environmental threats. It fills gaps, seals cracks, and minimizes heat loss during colder days. Propolis also exhibits antimicrobial properties, reducing infection risk and inhibiting the growth of fungi and bacteria, thereby creating a hygienic environment for bees (16). Propolis contains polyphenols, terpenoids, flavonoids, and aromatic acids, which are responsible for its strong antioxidant activity and help prevent the development of bacterial resistance (15,16). In addition to reducing bacterial mobility, propolis may disturb membrane potential, interfere with permeability of the microbe's cell membrane, and inhibit ATP generation (16). Few studies were conducted to compare the therapeutic effects of beehive products; therefore, in this study, we aimed to evaluate the antibacterial, antioxidant, and anti-inflammatory effects of beehive products (honey, propolis, and royal jelly) from two regions of Algeria.

METHODS

The study analyzed two samples each of honey, propolis, and royal jelly produced by *Apis mellifera* bees from a semi-arid region in the central part of Algeria and a humid region in the east of Algeria. The floral origin of the samples was determined based on the plant species present in the vicinity of each hive. All samples were kept at 6°C until their analysis. Honey and royal jelly were diluted using distilled water. For propolis, 1 g of propolis powder was homogenized and mixed with 10 mL of an 80% (v/v) ethanol/water solution for 72 hours and then kept at 70 °C for 1 hour. The resulting mixtures were filtered and evaporated. All samples were filtered through 0.22µm microfilters to eliminate bacterial contamination.

Table 1. Color, pH value, and floral and geographical origin of hive bee products

Samples	Color	pH	Floral origin	Region	Climate
Honey 1	Dark brown	3.15 ± 0.02	<i>Ziziphus sp, Brassica sp, Erica arborea, Carduus type, Lotus sp, Xanthium sp.</i>	Djelfa	Semi-arid
Propolis 1	Dark brown	4.19 ± 0.02			
Royal jelly 1	Cream	3.91 ± 0.02			
Honey 2	Brown	3.47 ± 0.01	<i>Trifolium sp, Eucalyptus sp, Rosmarinus sp, Hedera helix, Convolvulacea, Chrozophora tinctoria</i>	Annaba	Humid
Propolis 2	Dark brown	5.26 ± 0.03			
Royal jelly 2	Cream	3.24 ± 0.03			

The following dilutions were prepared: 2.5, 5, 10, 20, 40, 60, 80, 100% (v/v).

Antibacterial effect assays

The evaluation of the antibacterial activity of the samples was carried out according to the methods described by Bouacha M et al. towards ten multidrug-resistant bacteria isolated from infected wounds (8). These included Gram-negative bacteria: *Escherichia coli*, *Enterobacter aerogenes*, *Citrobacter koseri*, *Proteus mirabilis*, *Klebsiella pneumoniae*, *K. oxytoca*, and *Pseudomonas aeruginosa*; and Gram-positive bacteria: *Staphylococcus aureus*, *S. saprophyticus*, and *Enterobacter faecalis*. An inoculum of each bacterium was prepared by transferring 1 to 2 well-isolated and perfectly identical colonies into nutrient broth (Difco, MD, USA). The optical density was measured and adjusted to 0.08 and 0.10 at 625 nm.

Agar diffusion assay

The study used a standard technique to evaluate the efficacy of antimicrobial agents, specifically honey. Mueller Hinton agar (Difco, MD, USA) was prepared and poured into sterile Petri dishes, and wells of 6 mm in diameter were created as reservoirs for honey samples. Each plate was inoculated with a bacterial suspension standardized to an appropriate concentration. After inoculation, 50 µL of honey dilution was dispensed into each well, and the plates were incubated at 37°C for 24 hours. The antibacterial activity was assessed by measuring the diameter of clear zones of inhibition around the wells. The size of these zones correlated with the antibacterial potency of a honey sample, with larger zones indicating stronger antibacterial activity.

Microdilution assay

96-well microplates (Fisher Scientific, UK) were used to determine the minimum inhibitory concentration of honey (MIC) for antibacterial activity. For this test, 100 µL of bacterial inoculum was mixed in each well with 100 µL of honey at various dilutions. Wells containing inoculum but no honey were positive controls, while broth alone was negative. The microplates were incubated at 37°C for 24 hours to cultivate

bacteria.

Bacterial growth was evaluated by absorbance at 620 nm to derive MIC values. Formula for calculating growth inhibition: $\text{Inhibition (\%)} = \left(\frac{A_{\text{test}} - A_{\text{control}}}{A_{\text{test}}} \right) \times 100$, where A control and A test were the absorbance values of the control and test samples, respectively. The lowest honey dilution that inhibits bacterial growth 100%, showing no well growth, was the MIC. To determine the minimal bactericidal concentration (MBC), 10 µL from wells without growth was plated onto nutrient agar plates. These plates were incubated at 37°C for 24 hours. The MBC was defined as the lowest honey dilution at which no bacterial growth occurred, proving its bactericidal properties. To compare the honey's bacteriostatic (growth-inhibiting) and bactericidal (bacteria-killing) activities, the MBC/MIC ratio was calculated.

Antioxidant effect

Total polyphenolic content determination

The quantification of total polyphenols was carried out using the Folin-Ciocalteu assay (17). Briefly, a 5 mL aliquot of a methanolic hive product solution (1g in 5 mL) was mixed with 2.4 mL of distilled water and 200 µL of Folin-Ciocalteu reagent. After a 3-minute reaction period, 0.6 mL of a 20% sodium carbonate solution was added to the mixture. The samples were incubated in the dark at 25 °C for two hours. The absorbance of the reaction was measured at 725 nm using a spectrophotometer. Calibration was performed using various concentrations of gallic acid, and results were expressed in milligrams of gallic acid equivalents per 100 grams of honey (mg GAE/100g).

Flavonoids determination

Flavonoid content was determined using a colorimetric method involving aluminum chloride (18). A 1 mL sample of a 2% AlCl₃ solution was mixed with 1 mL of a methanolic solution of hive product (1 mg/mL). The mixture was allowed to react at 24 °C for 40 minutes. Absorbance readings were taken at 430 nm. A calibration curve was constructed using quercetin standards at concentrations of 20, 40, 60, 80, and 100 mg/L. Results were reported as milligrams of quercetin equivalents per gram of hive product (mg QE/g).

Ferric reducing/antioxidant power assay

The Ferric Reducing Antioxidant Power (FRAP) assay was performed following the method outlined by Beretta et al. (17). This colorimetric assay measures the change in absorbance at 593 nm, reflecting the reduction of colorless Fe³⁺-TPTZ (2,4,6-tripyridyl-s-triazine) to blue Fe²⁺-TPTZ by antioxidants able to donate electrons. Five products were prepared at a concentration of 1 g/mL in double-distilled water. A 200 µL aliquot of the honey solution was combined with 1.8 mL of FRAP reagent, consisting of 10 mmol/L TPTZ in 40 mmol/L HCl, 20 mmol/L FeCl₃, and 0.3 mol/L acetate buffer at pH 3.6. After incubating at 37 °C for 10 minutes, absorbance was measured at 593 nm. The antioxidant capacity was quantified in µmol/L of ascorbic acid equivalents per gram of honey (µmol/L AAE/g), using a standard curve prepared with ascorbic acid concentrations from 20 to 700 µmol/L.

1,1-Diphenyl-2-Picrylhydrazyl (DPPH) assay

The DPPH radical scavenging capacity was assessed using a modified method. A 2.7 mL solution of methanolic DPPH (6.10⁻⁵ mol/L) was mixed with 0.3 mL of honey. After vortexing the mixture at 2500 rpm for 1 minute, the solution was incubated in the dark for one hour. Following this incubation, the absorbance was measured at 517 nm after an additional 15-minute incubation at 25°C. Ascorbic acid served as the positive control. The DPPH radical scavenging activity (%) was calculated by taking the difference between the absorbance of the control and the sample, dividing it by the absorbance of the test, and then multiplying the result by 100. Using a calibration curve ranging from 0 to 10 mg/L of ascorbic acid, the IC₅₀ value, which represents the honey concentration required to scavenge 50% of DPPH radicals, was determined.

Anti-inflammatory effect

The albumin denaturation method, with a slight modification, was used to evaluate the anti-inflammatory activity *in vitro*, as described by Ali et al. (19). At varying concentrations, the samples were combined with a 1% water-based bovine serum albumin fraction in test tubes. Following a 15-minute incubation period at 37°C, the tubes were subjected to a 10-minute heating cycle at 70°C. Absorbance was measured spectrophotometrically at 660 nm. Distilled water was used as the negative control, while aspirin at the same concentrations (2.5%, 5%, 10%, 20%, 40%, 50%, 80%, and 100%) served as the positive control. The following formula was used to compute the percentage of inhibition of albumin denaturation = $\left(\frac{A_2 - A_1}{A_2}\right) \times 100$.

In this case, A1 represents honey's absorbance, and A2 represents the control's absorbance (distilled water).

Data analysis

Results were expressed as mean values ± standard deviation (SD), and the study used triple analyses to guarantee accuracy and reliability. The GraphPad Prism software was used to conduct the statistical analysis. To differentiate between treatments, a one-way analysis of variance (ANOVA) was employed. When comparing two sets of data, Tukey's post hoc test was used. Statistical analysis was performed with significance set at P < 0.05, supporting the reliability of the observed differences.

RESULTS

Antibacterial activity

As shown in Figure 1 and Table 2, the antibacterial activities of hive products have been thoroughly investigated. The results showed that the tested products varied significantly; honey samples outperformed royal jelly and propolis. Furthermore, the hive products had a greater impact on Gram-positive bacteria compared to Gram-negative bacteria. According to the MBC/MIC ratios, none of the bee products were bacteriostatic; rather, they were bactericidal. This indicates that the hive products could kill the bacteria, not just inhibit their growth. Based on these findings, honey, royal jelly, and propolis all demonstrated the ability to operate as natural antibacterial agents, with honey showing the highest efficacy.

Antioxidant effect

The determination of total phenolic content, flavonoid content, and antioxidant effects of hive products is reported in Figure 2. The total polyphenolic content of honey sample 1 was found to be relatively high in comparison with those of propolis and royal jelly. Flavonoid content also varied significantly; propolis collected from the Djelfa region has the highest concentration. Honey samples exhibited the highest antioxidant activity, followed by propolis and royal jelly samples.

Anti-inflammatory effect

The results of the anti-inflammatory activities of hive products determined by the membrane stabilization method are presented in Figure 3. The hive products showed a good anti-inflammatory activity. Honey and royal jelly samples were more effective than propolis.




Figure 1. Antibacterial effect of honey, propolis, and royal jelly using the well diffusion assay expressed as inhibitory diameters \pm SD (mm)

Table 2. MIC, MBC, and MBC/MIC ratio of honey, propolis, and royal jelly

	Honey 1		Honey 2		Propolis 1		Propolis 2		Royal jelly 1		Royal jelly 2	
	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC
<i>E. coli</i>	20.0	40	20	20	40	60	40	80	40	60	40	80
<i>E. aerogenes</i>	20.0	20	20	20	40	40	40	80	40	60	20	60
<i>C. koseri</i>	20.0	20	20	20	40	40	40	80	60	60	40	80
<i>P. mirabilis</i>	40.0	80	40	60	40	60	40	80	40	60	40	80
<i>K. pneumoniae</i>	40.0	100	40	60	40	100	40	100	40	100	40	100
<i>K. oxytoca</i>	40.0	60	40	80	40	80	40	80	40	100	40	100
<i>P. aeruginosa</i>	40.0	60	40	80	40	60	40	100	60	100	40	100
<i>S. aureus</i>	02.5	05.0	02.5	05.0	10	20	10	20	10	20	20	40
<i>S. saprophyticus</i>	02.5	05.0	02.5	05.0	5	10	10	20	10	20	20	40
<i>E. faecalis</i>	05.0	05.0	05.0	10.0	10	20	10	20	10	20	20	40
MBC/MIC ratio	01.0-02.5		01.0-02.0		01.0-2.0		01.0-02.5		01.0-02.5		01.0-03.0	




Figure 2. Antioxidant effects of honey, propolis, and royal jelly. (a): Total polyphenolic content, (b): flavonoid content, (c): DPPH, (d): FRAP. There are significant differences between samples with different letters.




Figure 3. The anti-inflammatory effects of honey, propolis, and royal jelly.

DISCUSSION

Hive products showed diversity in their bioactive components, contributing to their effectiveness against a variety of microorganisms, including multidrug-resistant bacteria. As reported in Figure 1 and Table 2, the antibacterial effect of honeybee products varied significantly. Honey samples showed more efficiency than the other products. Similarly to this result, (20) have also shown that honey exhibited the highest antibacterial activity against *E. coli* and *S. typhimurium* strains. The effectiveness of honey as an antibacterial agent can be attributed to several factors, such as acidity, glucose oxidase, and lysosomes, which may act in a synergetic manner to inhibit the growth and viability of pathogenic microorganisms (10). This finding was previously reported by many authors (2,3,21). The mechanism of action of the antibacterial effect of hive products is not well known. However, some authors suggest that they cause a disruption of membrane potential and permeability, which may contribute significantly to overall cytotoxicity, affecting bacterial viability. In addition, Gram-positive bacteria were more susceptible to hive products than Gram-negative bacteria. This is due to the structure of the outer membrane of Gram-negative bacteria. Indeed, Gram-positive and Gram-negative bacteria have a cell wall that can protect the cytoplasmic membrane. In addition, Gram-negative bacteria have additional protection offered by the polysaccharide-rich outer membrane (11). Moreover, the tested hive products exhibit a bactericidal effect on both Gram-negative and Gram-positive bacteria, as their MBC/MIC ratios are between 1 and 3. This ratio is essential for differentiating between a bacteriostatic agent, which prevents bacterial proliferation without killing the bacterium, and a bactericidal agent, which destroys and kills the bacterial cell. According to O'Neill and Chopra

(22), an antimicrobial agent is bactericidal when the MBC/MIC ratio is less than or equal to 4; therefore, the tested hive products exhibited a bactericidal effect against multidrug-resistant bacteria (22). Many authors previously reported that honey (3,11,23,24) and propolis (25–27) displayed a bactericidal effect. However, García MC et al. (14) reported that two royal jelly samples had a bacteriostatic effect.

The determination of the total polyphenolic content, flavonoid content, and antioxidant effects of hive product samples (Figure 2) showed significant differences among them. Propolis had the highest TPC and TFC content when compared with honey and royal jelly. However, honey had the highest antioxidant activity, suggesting that it may contain other compounds with antioxidant effects that can protect against oxidative stress. Similarly, propolis and royal jelly have also been found to have antioxidant properties due to their high phenolic and flavonoid content. Several mechanisms contribute to the antioxidant effects of hive products, including free-radical scavenging, hydrogen donation, metal ion chelation, quenching of singlet oxygen, and acting as a substrate for superoxide and hydroxyl radicals. According to many authors, phenolic compounds are strongly correlated with antioxidant activity of bee products, which is influenced by botanical source, geographic and entomological origin, and climatic conditions. (3,11,28,29). However, hive products contain other substances with antioxidant effects, such as minerals, amino acids, peptides, proteins, organic acids, and enzymes (15,30). Similar results were previously reported by Nagai T et al. (31). In this study, commercial honeys, royal jelly, and propolis were found to have a significant antioxidant activity, as measured by their ability to scavenge free radicals and inhibit lipid peroxidation (32). Mouhoubi-Tafinine et al. also found that honey samples had a higher antioxidant activity than propolis samples (12). In contrast, Postali E et al. reported that propolis showed the highest antioxidant activity (33). Buratti S et al. also found that propolis samples had a higher antioxidant capacity than honey (34). Indeed, many factors could affect the similarity or divergence in the antioxidant effect of hive products, including bee species, geographic region, plant species, forage, harvest, and storage conditions. Additionally, the methods used to evaluate antioxidant activity can significantly impact results. Variations in sample dilution, extraction methods, and conditions can affect the concentration of bioactive compounds and their potency. Criteria for reporting findings can also cause discrepancies in results.

Therefore, a standardized approach is crucial for reliable and comparable results in antioxidant activity evaluations. Based on the results in Figure 3, all hive products exhibited the anti-inflammatory effect; however, honey samples were found to have the most potent anti-inflammatory effects, followed by royal jelly and propolis. The anti-inflammatory activity of honey is attributed to its capacity to block the synthesis of pro-inflammatory cytokines, such as tumor necrosis factor-alpha (TNF- α), interleukin-6 (IL-6), and interleukin-1 beta (IL-1 β) (35). The antioxidant and anti-inflammatory properties of hive products are closely connected and can be interdependent. The qualities are mostly attributable to the presence of polyphenols and flavonoids, which can modify the immune response, decrease pro-inflammatory cytokine production, scavenge free radicals, and diminish oxidative stress (15,21). The study demonstrated that honey, propolis, and royal jelly have significant bactericidal effects against a variety of microorganisms, including multidrug-resistant bacteria. All hive products showed good antioxidant and anti-inflammatory properties due to their phenolic and flavonoid content. Honey samples exhibited the strongest antibacterial, antioxidant, and anti-inflammatory properties. Overall, bee products have potential as natural alternatives for various health conditions. However, further research is needed to better understand their mechanisms of action and to establish optimal dosages and formulations for therapeutic use.

REFERENCES

1. Lannutti L, Gonzales FN, Dus Santos MJ, Florin-Christensen M, Schnittger L. Molecular detection and differentiation of arthropod, fungal, protozoan, bacterial and viral pathogens of honeybees. *Vet Sci*. 2022;9(5):221. [\[CrossRef\]](#)
2. Bouacha M, Ayed H, Grara N. Honey bee as alternative medicine to treat eleven multidrug-resistant bacteria causing urinary tract infection during pregnancy. *Sci Pharm*. 2018;86(2):14. [\[CrossRef\]](#)
3. Boudiar I, Bouacha M, Besnaci S, Khallef M, Abdi A. Evaluation of the antibacterial and antimutagenic effects of multiflora honeys produced by the *Apis mellifera* bee and their correlation with polyphenolic content, flavonoids and color. *J Entomol Res*. 2022;46(3):607-14. [\[CrossRef\]](#)
4. Bouacha M, Boudiar I, Akila A, Al-Kafawween MA, Khallef M. The antimutagenic effect of multiflora honey in *Salmonella*/microsomal assay and its correlation with the total polyphenolic content. *J Microbiol Biotechnol Food Sci*. 2022;11(6):e5557. [\[CrossRef\]](#)
5. Ahmed S, Sulaiman SA, Baig AA, et al. Honey as a potential natural antioxidant medicine: an insight into its molecular mechanisms of action. *Oxid Med Cell Longev*. 2018; 2018:8367846. [\[CrossRef\]](#)
6. Abbas M, Shah I, Nawaz MA, et al. Honey against cancer. In: Jafari SM, Nabavi SM, Silva AS, editors. *Nutraceuticals: Cancer Signaling and Clinical Aspects of Mode of Action*. Cham: Springer; 2021; 401-8. [\[CrossRef\]](#)
7. Papa G, Maier R, Durazzo A, et al. The honey bee *Apis mellifera*: an insect at the interface between human and ecosystem health. *Biology*. 2022;11(2):233. [\[CrossRef\]](#)

Acknowledgements

This study was not supported by any sponsor or funder.

Authors' Contributions

Conceptualization, M.B. and I.B.; Methodology, M.B. and I.B.; Investigation, M.B.; Formal Analysis, I.B.; Writing – original draft, M.B.; Writing – review & editing, M.B. and I.B. Both authors have read and approved the published version of the manuscript.

Statement of Competing Interest

The authors declare no relevant conflicts of interest.

Statement of Data Availability

The datasets generated and analyzed during the current study are available from the corresponding author.

Statement of Generative AI Use

No generative AI was used.

Publisher's Note: The statements, opinions, and data contained in AFMN Biomedicine articles are solely those of the individual author(s) and contributor(s) and do not necessarily represent the views of the publisher or the editor(s). The publisher and editor(s) disclaim responsibility for any harm or damage caused by the use of information or products mentioned in the publication.

8. Bouacha M, Besnaci S, Boudiar I. An overview of the most used methods to determine the in vitro antibacterial activity of honey. *Acta Microbiol Bulg.* 2023;39(1):23-30. [[CrossRef](#)]
9. Bouacha M, Besnaci S, Boudiar I. Screening of the antibacterial and anti-biofilm effect of multifloral honey against multidrug-resistant *Pseudomonas aeruginosa*. *Acta Microbiol Hell.* 2022;67(1).
10. Molan PC. The antibacterial activity of honey: 2. Variation in the potency of the antibacterial activity. *Bee World.* 1992;73(2):59-76. [[CrossRef](#)]
11. Bouacha M, Besnaci S, Boudiar I, Al-Kafawween MA. Impact of storage on honey antibacterial and antioxidant activities and their correlation with polyphenolic content. *Trop J Nat Prod Res.* 2022;6(1):34-9. [[CrossRef](#)]
12. Al-Mamary M, Al-Meer A, Al-Habori M. Antioxidant activities and total phenolics of different types of honey. *Nutr Res.* 2002;22(9):1041-7. [[CrossRef](#)]
13. Ranneh Y, Akim AM, Hamid HA, et al. Honey and its nutritional and anti-inflammatory value. *BMC Complement Med Ther.* 2021;21(1):17. [[CrossRef](#)]
14. García MC, Finola MS, Marioli JM. Antibacterial activity of royal jelly against bacteria capable of infecting cutaneous wounds. *J ApiProd ApiMed Sci.* 2010;2(3):93-9. [[CrossRef](#)]
15. Viuda-Martos M, Ruiz-Navajas Y, Fernández-López J, Pérez-Álvarez JA. Functional properties of honey, propolis, and royal jelly. *J Food Sci.* 2008;73(9):R117-R124. [[CrossRef](#)]
16. Przybyłek I, Karpiński TM. Antibacterial properties of propolis. *Molecules.* 2019;24(11):2047. [[CrossRef](#)]
17. Beretta G, Granata P, Ferrero M, Orioli M, Maffei Facino R. Standardization of antioxidant properties of honey by a combination of spectrophotometric/fluorimetric assays and chemometrics. *Anal Chim Acta.* 2005;533(2):185-91. [[CrossRef](#)]
18. Meda A, Lamien CE, Romito M, Millogo J, Nacoulma OG. Determination of the total phenolic, flavonoid and proline contents in Burkina Fasan honey, as well as their radical scavenging activity. *Food Chem.* 2005;91(3):571-7. [[CrossRef](#)]
19. Ali SS, Morsy R, El-Zawawy NA, Fareed MF, Bedaiwy MY. Synthesized zinc peroxide nanoparticles (ZnO₂-NPs): a novel antimicrobial, anti-elastase, anti-keratinase, and anti-inflammatory approach toward polymicrobial burn wounds. *Int J Nanomedicine.* 2017;12:6059-78. [[CrossRef](#)]
20. Badawy OFH, Shafii SSA, Tharwat EE, Kamal AM. Antibacterial activity of bee honey and its therapeutic usefulness against *Escherichia coli* O157:H7 and *Salmonella typhimurium* infection. *Rev Sci Tech.* 2004;23(3):1011-22. [[CrossRef](#)]
21. Nader RA, Mackieh R, Wehbe R, El Obeid D, Sabatier JM, Fajloun Z. Beehive products as antibacterial agents: a review. *Antibiotics.* 2021;10(6):717. [[CrossRef](#)]
22. O'Neill AJ, Chopra I. Preclinical evaluation of novel antibacterial agents by microbiological and molecular techniques. *Expert Opin Investig Drugs.* 2004;13(8):1045-63. [[CrossRef](#)]
23. Bazaid AS, Aldarhami A, Patel M, et al. The antimicrobial effects of Saudi Sumra honey against drug resistant pathogens: phytochemical analysis, antibiofilm, anti-quorum sensing and antioxidant activities. *Pharmaceuticals.* 2022;15(10):1212. [[CrossRef](#)]
24. Yu L, Palafox-Rosas R, Luna B, She RC. The bactericidal activity and spore inhibition effect of Manuka honey against *Clostridioides difficile*. *Antibiotics.* 2020;9(10):684. [[CrossRef](#)]
25. de Souza Silva T, Silva JM, Braun GH, et al. Green and red Brazilian propolis: antimicrobial potential and anti-virulence against ATCC and clinically isolated multidrug-resistant bacteria. *Chem Biodivers.* 2021;18(8):e2100307. [[CrossRef](#)]
26. Özüşağlam MA, Tacer S, Boulechfar S, Zellagui A. An investigation of the bactericidal and fungicidal effects of Algerian propolis extracts and essential oils. *Cumhuriyet Sci J.* 2022;43(1):14-9. [[CrossRef](#)]
27. Widelski J, Okirčyc P, Suśniak K, et al. Correlation between chemical profile of Georgian propolis extracts and their activity against *Helicobacter pylori*. *Molecules.* 2023;28(3):1374. [[CrossRef](#)]
28. Al-Farsi M, Al-Amri A, Al-Hadhrami A, Al-Belushi S. Color, flavonoids, phenolics and antioxidants of Omani honey. *Heliyon.* 2018;4(10):e00874. [[CrossRef](#)]
29. Can Z, Yildiz O, Sahin H, Turumtay EA, Silici S, Kolayli S. An investigation of Turkish honeys: their physico-chemical properties, antioxidant capacities and phenolic profiles. *Food Chem.* 2015;180:133-41. [[CrossRef](#)]
30. Martinello M, Mutinelli F. Antioxidant activity in bee products: a review. *Antioxidants.* 2021;10(1):71. [[CrossRef](#)]
31. Nagai T, Sakai M, Inoue R, Inoue H, Suzuki N. Antioxidative activities of some commercially available honeys, royal jelly, and propolis. *Food Chem.* 2001;75(2):237-40. [[CrossRef](#)]
32. Mouhoubi-Tafinine Z, Ouchemoukh S, Tamendjari A. Antioxidant activity of some Algerian honey and propolis. *Ind Crops Prod.* 2016;88:85-90. [[CrossRef](#)]
33. Postali E, Peroukidou P, Giaouris E, Papachristoforou A. Investigating possible synergism in the antioxidant and antibacterial actions of honey and propolis from the Greek island of Samothrace through their combined application. *Foods.* 2022;11(14):2041. [[CrossRef](#)]
34. Buratti S, Benedetti S, Cosio MS. Evaluation of the antioxidant power of honey, propolis and royal jelly by amperometric flow injection analysis. *Talanta.* 2007;71(3):1387-92. [[CrossRef](#)]
35. Iftikhar A, Nausheen R, Mukhtar I, et al. The regenerative potential of honey: A comprehensive literature review. *J Apic Res.* 2023;62(1):97-112. [[CrossRef](#)]