

VACCINE HESITANCY IN PARENTS OF CHILDREN WITH CONGENITAL UROGENITAL ANOMALIES WITH HYPOSPADIAS AND UNDESCENDED TESTIS

Mehmet Semih Demirtaş¹  Siddika Songül Yalçın²  Mustafa Tuşat³ 
Ismail Özmen⁴  Dilek Başar⁵  Ahmet Salih Calapoglu⁶ 

¹Department of Pediatrics, Division of Social Pediatrics, Aksaray University Faculty of Medicine, Aksaray, Turkey ²Department of Social Pediatrics, Hacettepe University Faculty of Medicine, Ankara, Turkey ³Department of Pediatric Surgery, Aksaray University Faculty of Medicine, Aksaray, Turkey ⁴Department of Pediatric Surgery, Kilis Prof. Dr. Alaeddin Yavasca State Hospital, Kilis, Turkey ⁵Department of Pediatric Surgery, Trabzon Kanuni Training and Research Hospital, Trabzon, Turkey ⁶Department of Pediatric Surgery, Recep Tayyip Erdoğan University Faculty of Medicine, Rize, Turkey

Parental vaccine hesitancy (VH) toward childhood immunization is a critical public health challenge affecting individual pediatric health and broader societal immunity. This study aimed to evaluate VH among parents of male children by comparing those whose children had congenital urogenital anomalies (CUA), specifically hypospadias and undescended testicles, with those of children without malformations. The assessment also incorporated family demographics, health literacy levels, and adherence to preventive health practices. This descriptive-analytical study included 409 participants, divided into two groups: parents of children presenting for elective circumcision without malformations (CNM) and parents of children diagnosed with CUA. Data were collected using a socio-demographic form, the Turkey Health Literacy Scale (TSOY-32), and the Vaccine Hesitancy Scale (VHS). High vaccine hesitancy was defined as scores in the upper quartile of the VHS (VHS-uQ). The overall vaccination rate among the children was 96.6% (n = 395). In the CUA group, the absence of routine vitamin D supplementation was associated with 25-fold higher odds of VHS-uQ (p < 0.001). Mothers with a high school education or higher demonstrated a 4.69-fold higher odds of VHS-uQ (p < 0.001). Conversely, COVID-19 vaccination of both parents was associated with lower VHS-uQ rates in both groups (p < 0.001). Interestingly, higher TSOY-32 scores were associated with a 4.57-fold higher odds of VHS-uQ (p = 0.034). This study identified maternal health literacy and adherence to routine preventive care as key determinants of VH among parents of children with CUA. Findings suggest that fostering trust through continuous healthcare engagement and evidence-based counseling is a vital pathway to improving vaccine acceptance in this specific population.

Keywords: vaccine hesitancy, childhood vaccination, parental vaccine hesitancy, hypospadias, undescended testicles, congenital urogenital anomalies

Submitted: February 7, 2025 **Accepted:** June 18, 2025

Published online: June 15, 2026

Copyright: © 2026, Author(s). This is an open-access article published under the terms of the Creative Commons Attribution 4.0 International License. (<http://creativecommons.org/licenses/by/4.0/>).

Correspondence to:

Mehmet Semih Demirtaş
Department of Pediatrics Division of Social Pediatrics Centre
Aksaray University Faculty of Medicine
Aksaray, Turkey
E-mail: md.semihdemirtas@gmail.com

INTRODUCTION

Hypospadias and undescended testes represent the most prevalent congenital urogenital anomalies (CUA) in male children (1,2). The diagnosis of a congenital anomaly, combined with the necessity for surgical intervention and the potential for complications, often induces significant anxiety and fear in parents. Such emotional stressors can profoundly influence the healthcare-related decisions parents make for their children (3,4). Parents typically acquire health information from mass media, their social environment, and healthcare professionals to manage their children's daily care and long-term well-being (5,6). Within the spectrum of pediatric healthcare, vaccination remains a fundamental and highly effective primary public health strategy for disease prevention (7).

Substantial progress has been achieved through routine childhood immunization, which has targeted infections that were the primary causes of pediatric morbidity and mortality in the last century. It is estimated that global vaccination programs, which have led to the eradication of smallpox and elimination of polio in numerous regions, save more than 3 million children annually (8). However, vaccine hesitancy (VH), identified by the World Health Organization (WHO) in 2019 as one of the top ten global health threats, has increased gradually. This trend has intensified following the COVID-19 pandemic, incited by social media misinformation regarding both COVID-19 and routine childhood vaccines (9,10). VH among parents striving to make optimal health decisions for their children poses a significant challenge, impacting not only individual pediatric health but also collective societal immunity. Evaluating determinants of VH, such as sociocultural influences, economic conditions, environmental factors, and media exposure, is essential for providing timely, evidence-based guidance to parents (9,11,12).

In this study, we aimed to evaluate and compare VH levels among parents of male children diagnosed with CUA (hypospadias and undescended testes) and those of children presenting for elective circumcision without malformations (CNM).

METHODS

Study design and settings

This multicenter, descriptive-analytical study was conducted between February 2024 and May 2024 across four consortium centers in Turkey (Aksaray, Rize, Trabzon, and Kilis). These centers represent three distinct regions,

according to the Nomenclature of Territorial Units for Statistics (NUTS-1) and the Turkey Demographic and Health Survey (TDHS) (13).

Study population and sampling

A total of 492 parents of male children aged 0–7 years, who presented to pediatric surgery outpatient clinics for hypospadias, undescended testes, or elective circumcision, were initially recruited using stratified sampling. Participants were categorized into two groups: those with CUA and those seeking CNM.

Exclusion criteria were as follows: 35 participants with comorbidities, 9 who provided uniform (identical) answers to all scale items, 33 patients aged 98 months or older, and 6 who provided contradictory responses. Consequently, the final analysis was performed on 409 participants.

Data collection and instruments

Data were collected through face-to-face interviews conducted by pediatric surgeons at the respective centers. The research instrument consisted of a 31-item structured questionnaire (Supplementary File 1), the Turkey Health Literacy Scale (TSOY-32) (14), and the Vaccine Hesitancy Scale (VHS) (15).

The structured questionnaire comprised five sections:

1. Sociodemographic information: parental education levels, ages, and child-specific data (birth order, weight, and gestational age).
2. Healthcare practices: adherence to routine infant-child-adolescent monitoring protocols of the Turkish Ministry of Health (16), including iron/vitamin D supplementation, antenatal care visits, and maternal vaccination status.
3. Vaccination history and attitudes: perspectives on routine and private vaccines (e.g., rotavirus, meningococci), COVID-19 vaccination status, and the impact of social media on vaccine perception (17-19).
4. Information sources: preferred sources for health information (e.g., pediatricians, family physicians, social media, bloggers, and official websites).

TSOY-32 scale

Maternal health literacy (HL) was assessed using the TSOY-32 scale. This 32-item, 5-point Likert-type scale evaluates four dimensions—accessing, understanding, evaluating, and applying health information—across two domains: "Treatment and Service Utilization" and "Disease Prevention/Health Promotion" (20,21). For the score

calculation of the scale consisting of 32 items on a 5-point Likert scale (1=Very easy, 2=Easy, 3=Difficult, 4=Very difficult, 5=No idea) for each item, it was coded as suggested in the literature, and a value of 0 was assigned to the 'no idea' code. Scores ranging from 0 (lowest HL) to 50 (highest HL) were categorized into three levels: 0–25, >25–33, and >33–50 points.

Vaccine hesitancy scale

The Turkish version of the 9-item VHS was administered to assess parental hesitancy (15). Items are scored on a 5-point Likert scale, with total scores ranging from 9 to 45; higher scores indicate greater hesitancy. For analysis, participants were dichotomized into two groups based on the upper quartile: lower quartile (VHS-LQ; <25 points) and upper quartile (VHS-UQ; ≥25 points).

Ethics

The study was conducted in accordance with the Declaration of Helsinki and was approved by the Hacettepe University non-interventional Ethics Committee (Approval No. GO 2024/05-42) on January 23, 2024.

Statistical analyses

In the study, the data were collected in Excel format, and analyses were performed using SPSS 22.0 (IBM, USA).

Participants were divided into two groups: those with congenital urogenital anomalies (CUA) and the CNM group.

VHS (Vaccine Hesitancy Scale) scores were evaluated in quartiles, divided into the upper quartile (VHS-UQ) (Q4) and lower quartiles (Q1-3) (VHS-LQ).

The normal distribution of the data was evaluated by the Shapiro-Wilk test. Categorical variables were expressed as n (%) and continuous variables showing normal distribution were expressed as mean ± standard deviation. The Chi-square test was used to compare the percentage distribution of categorical data between groups.

Variables with an association to VH indicated by a p-value of <0.10 were selected for further analysis. Multiple logistic regression analysis was then used to explore associations between child-family parameters (such as sociodemographic characteristics, vaccination practices, healthcare practices, and health information sources) and the VHS-UQ in both the CUA and CNM groups. Additionally, data from the overall group were analyzed to identify associations between VH and child-family parameters,

including TSOY-32 scale scores and the different study groups. Adjusted odds ratios with 95% confidence intervals were calculated, and a p-value of <0.05 was considered statistically significant.

RESULTS

Sociodemographic characteristics

In this study, 36.7% (n = 150) of the children were first-borns. Regarding age distribution, 48.9% were younger than 24 months, 29.3% were aged 24–47 months, and 21.8% were 48 months or older. Routine vitamin D supplementation was administered to 90.7% of the participants, while 74.4% adhered to the standard iron supplementation regimen starting at 4 months of age (Table 1).

Maternal age was distributed as follows: 10.3% were under 25 years, 65.3% were between 25 and 34 years, and 24.4% were 35 years or older. In contrast, 55% of the fathers were under 25 years old. The sociodemographic profiles of the CUA and CNM groups were largely comparable, with no significant differences in maternal age (p = 0.585), paternal age (p = 0.544), gestational duration (p = 0.085), child age distribution (p = 0.264), or birth weight (p = 0.232). However, maternal education levels differed significantly: 54.2% of mothers in the CUA group had eight years or less of education compared to 41.9% in the CNM group (p = 0.015) (Table 1).

Vaccination rates and attitudes

The overall childhood vaccination rate was 96.6% (n = 395). While the CUA group showed a slightly higher routine vaccination rate (98%) compared to the CNM group (94.4%), this difference was at the threshold of significance (p = 0.050). Parental COVID-19 vaccination rates were similar between groups (CUA fathers: 61.0% vs. CNM fathers: 63.7%, p = 0.582; overall mothers: 66.5%).

Post-pandemic attitudes revealed that while 37.7% of parents viewed routine childhood vaccines as more essential after COVID-19, a substantial minority (24.9% overall) reported increased hesitancy. Negative impacts from social media news regarding vaccines were reported by 16.4% of parents, with a higher but not statistically significant prevalence in the CUA group (18.5% vs. 13.1%, p = 0.215). Uptake for special vaccines (rotavirus/meningococcal) remained low at 12.5% across both groups (p = 0.550).

Table 1. Evaluation of socio-demographic, health practice, and vaccination information of parents

Features	Overall n (%)	CUA* (n = 249)	Control (n = 160)	p/E
Child characteristics				
First child				
1st	150 (36.7)	86 (34.5)	64 (40.0)	0.263
2nd and more	259 (63.3)	163 (65.5)	96 (60.0)	
Child age (month)				
<24	200 (48.9)	114 (45.8)	86 (53.8)	0.264
24-47	120 (29.3)	76 (30.5)	44 (27.5)	
≥48	89 (21.8)	59 (23.7)	30 (18.8)	
Gestational duration				
Mature	339 (82.9)	213 (85.5)	146 (91.3)	0.085
Premature	70 (17.1)	36 (14.5)	14 (8.8)	
Birth weight				
SGA3	50 (12.2)	18 (14.4)	16 (9.3)	0.232
No	359 (87.8)	122 (85.6)	170 (90.7)	
Hospitalization				
Yes	112 (27.4)	74 (29.7)	38 (23.8)	0.186
No	297 (72.6)	175 (70.3)	122 (76.3)	
Childhood vaccination§				
Yes	395 (96.6)	244 (98.0)	151 (94.4)	0.05
No	14 (3.4)	5 (2.0)	9 (5.6)	
At least 1 dose of special vaccine&				
Yes	51 (12.5)	33 (13.3)	18 (11.3)	0.55
No	358 (87.5)	216 (86.7)	142 (88.8)	
Parent characteristics				
Maternal age				
<25	42 (10.3)	23 (9.2)	19 (11.9)	0.585
25-34	267 (65.3)	162 (65.1)	105 (65.6)	
≥35	100 (24.4)	64 (25.7)	36 (22.5)	
Paternal age				
<35	225 (55.0)	134 (53.8)	91 (56.9)	0.544
≥35	184 (45.0)	115 (46.2)	69 (43.1)	
Maternal education				
8 years and lessß	202 (49.4)	135 (54.2)	67 (41.9)	0.015
High school and above	207 (50.6)	134 (53.8)	93 (58.1)	
Paternal education				
8 years and lessß	186 (45.6)	115 (46.2)	71 (44.4)	0.72
High school and above	223 (54.5)	79 (63.2)	89 (55.6)	
Health care practices				
Antenatal care visit				
1-3 times	42 (10.5)	29 (11.8)	13 (8.3)	0.264
4 and more	359 (89.5)	216 (88.2)	143 (91.7)	
Vaccination during pregnancy				
Yes	308 (75.3)	182 (73.1)	126 (78.8)	0.195
No	101 (24.7)	67 (26.9)	34 (21.3)	
Standard vitamin D supplementation regimen during infancy				
Yes	371 (90.7)	228 (91.2)	144 (90.0)	0.692
No	38 (9.3)	22 (8.8)	16 (10.0)	
Standard iron supplementation regimen during infancy				
Yes	300 (73.3)	172 (69.1)	128 (80.0)	0.015
No	109 (26.7)	77 (30.9)	32 (20.0)	

Standard vitamin D supplementation regimen during pregnancy				
Yes	365 (89.2)	226 (90.8)	139 (86.9)	0.216
No	44 (10.8)	23 (9.2)	21 (13.1)	
Standard iron supplementation regimen during pregnancy				
Yes	327 (80.0)	201 (80.7)	126 (78.8)	0.627
No	82 (20.0)	48 (19.3)	34 (21.3)	
Vaccine history and ideas of parent				
Childhood vaccination\$ idea				
Protects the child	313 (76.5)	187 (75.1)	126 (78.8)	0.447
I'm hesitant ☹	77 (18.8)	50 (20.1)	27 (16.9)	
Not safe ☹	12 (2.9)	9 (3.6)	3 (1.9)	
Refuse	7 (1.7)	3 (1.2)	4 (2.5)	
Childhood vaccination\$ idea after COVID-19				
Yes, necessary	154 (37.7)	93 (37.3)	61 (38.1)	0.811
Yes, hesitant	102 (24.9)	60 (24.1)	42 (26.3)	
No	153 (37.4)	96 (38.6)	57 (35.6)	
COVID-19 vaccination				
Mother	66 (16.1)	45 (18.1)	21 (13.1)	0.454
Father	48 (11.7)	26 (10.4)	22 (13.8)	
Parent	206 (50.4)	126 (50.6)	80 (50.0)	
No	89 (21.8)	52 (20.9)	37 (23.1)	
Maternal COVID-19 vaccination				
Yes	272 (66.5)	171 (68.7)	101 (63.1)	0.246
No	137 (33.5)	78 (31.3)	59 (36.9)	
Paternal COVID-19 vaccination				
Yes	254 (62.1)	152 (61.0)	102 (63.7)	0.582
No	155 (37.9)	97 (39.0)	58 (36.3)	
Influenced by social media news about vaccines				
Yes, hesitant	114 (27.9)	72 (28.9)	42 (26.3)	0.215
Yes, negative	67 (16.4)	46 (18.5)	21 (13.1)	
No, not affect	228 (55.7)	131 (52.6)	97 (60.6)	
Health information sources				
Family elders as a SHI†				
Reliable	274 (67.0)	164 (65.9)	110 (68.8)	0.545
Others	135 (33.0)	85 (34.1)	50 (31.3)	
Pediatrician as a SHI†				
Reliable	400 (97.8)	241 (96.8)	159 (99.4)	0.082
Others	9 (2.2)	8 (3.2)	1 (0.6)	
Family physician as a SHI†				
Reliable	382 (93.4)	232 (93.2)	150 (93.8)	0.819
Others	27 (6.6)	17 (6.8)	10 (6.3)	
Family nurse/midwife as a SHI†				
Reliable	352 (86.1)	215 (86.3)	137 (85.6)	0.837
Others	57 (13.9)	34 (13.7)	23 (14.4)	
Both members of the family medicine unit as a SHI†				
Reliable	344 (84.1)	208 (83.5)	136 (85.0)	0.5
One of them is reliable	46 (11.2)	31 (12.4)	15 (9.4)	
Others	19 (4.6)	10 (4.0)	9 (5.6)	
Social media as a SHI†				
Reliable	18 (4.4)	11 (4.4)	7 (4.4)	0.984
Others	391 (95.6)	238 (95.6)	153 (95.6)	

Government and university website as a SHI†				
Reliable	87 (21.3)	47 (18.9)	40 (25.0)	0.14
Others	322 (78.7)	202 (81.1)	120 (75.0)	
Social media blogger as a SHI†				
Reliable	34 (8.3)	22 (8.8)	12 (7.5)	0.633
Others	375 (91.7)	227 (91.2)	148 (92.5)	
Book as a SHI†				
Reliable	197 (48.2)	118 (47.4)	79 (49.4)	0.695
Others	212 (51.8)	131 (52.6)	81 (50.6)	
Television as a SHI†				
Reliable	46 (11.2)	31 (12.4)	15 (9.4)	0.337
Others	363 (88.8)	218 (87.6)	145 (90.6)	
Scales				
VHS Score π				
Lower Qα	306 (74.8)	182 (73.1)	124 (77.5)	0.316
Upper Qα	103 (25.2)	67 (26.9)	36 (22.5)	
TSOY**				
0-25	84 (43.3)	34 (39.1)	50 (46.7)	0.042
>25-33	58 (29.9)	22 (25.3)	36 (33.6)	
>33	52 (26.8)	31 (35.6)	21 (19.6)	

*=Congenital urogenital anomaly, †= Chi-square test was used to compare the CUA and control group. B= Primary and secondary education (8 years), §=Childhood routine vaccination programme, †= source of health information, ‡SGA= Small gestational age, §=Childhood routine vaccination programme, π= Vaccine hesitancy score, α= Lower Q group: <25 points; Upper Q group: ≥25 points &=Rota or meningococcal vaccine, ∅=I'm hesitant, but there's a vaccine tracking system. Θ=Not safe, but there's a vaccine tracking system, †= Source of health information, **= Turkish Health Literacy Scale

Factors associated with the vaccine hesitancy scale (VHS) Vaccine hesitancy (VHS-uQ) was significantly lower in children under 24 months (15.5%) compared to older age groups ($p < 0.001$). Furthermore, while only 14.4% of parents who expressed confidence in the national immunization program were in the VHS-uQ group, this rate rose to 59.7% among those who were hesitant and 91.7% among those who perceived vaccines as unsafe ($p < 0.001$) (Table 2). In the CUA group, maternal education level was a significant predictor: 36.8% of mothers with at least a high school education were in the VHS-uQ group compared to 18.5% of those with lower education levels ($p = 0.001$). Additionally, higher VHS scores were significantly associated with non-adherence to routine preventive care, including lack of routine vitamin D (72.7%) or iron supplementation (69.1%) during infancy ($p < 0.001$ and $p = 0.004$, respectively), and absence of maternal prenatal supplementation ($p < 0.001$). Distrust in the family medicine team was also linked to higher hesitancy ($p = 0.001$) (Table 2). In the CNM group, 45.2% of those who developed vaccine hesitancy post-pandemic fell into the VHS-uQ category ($p < 0.001$). Parents who were not vaccinated against COVID-19 showed significantly higher hesitancy rates (48.6%) compared to their vaccinated counterparts (11.3%, $p < 0.001$) (Table 2).

Logistic regression analysis Multivariable analysis of the entire cohort showed no significant association between VH and urogenital anomaly status or basic healthcare practices (all $p > 0.05$). However, significantly higher odds of VH were observed among participants with higher TSOY-32 scores (OR = 4.57, 95% CI: 1.12–18.65) and parents of children aged ≥48 months (OR = 4.13, 95% CI: 1.32–12.97). Conversely, parental COVID-19 vaccination was a protective factor against VH (Table 3). In the CUA group, the odds of high vaccine hesitancy were 6.31 times higher among mothers with at least a high school education ($p < 0.001$). Lack of routine vitamin D supplementation for the infant was associated with a 25-fold increase in the odds of VH (95% CI: 4.04–156.5). Vaccination of both parents against COVID-19 was strongly associated with lower VH scores (OR: 0.07, 95% CI: 0.02–0.24). In the CNM group, paternal education (high school or above) was associated with 7.52 times higher odds of VH ($p < 0.001$). Parents who reported increased hesitancy specifically after the COVID-19 pandemic had 11.43 times higher odds of being in the VHS-uQ group (95% CI: 2.52–51.84, $p = 0.002$) (Table 4).

Table 2. Evaluation of parents' vaccination hesitancy using the Vaccine Hesitancy Scale

Features	All			CUA			Control		
	Overall n = 409	VHS π Upper Q α		Overall Cua n = 249	VHS π Upper Q α		Overall Control n = 160	VHS π Upper Q α	
	n (%)	%**	p	n (%)	%**	p	n (%)	%**	P
Child characteristics									
First child			0.958			0.52			0.315
1st	150 (36.7)	25.3		86 (34.5)	24.4		64 (40.0)	26.6	
2nd and more	259 (63.3)	25.1		163 (65.5)	28.2		96 (60.0)	19.8	
Child age (month)			<0.001			0.003			0.02
<24	200 (48.9)	15.5		114 (45.8)	16.7		86 (53.8)	14	
24-47	120 (29.3)	32.5		76 (30.5)	32.9		44 (27.5)	31.8	
≥48	89 (21.8)	37.1		59 (23.7)	26.9		30 (18.7)	33.3	
Gestational duration			0.678			0.553			0.139
Mature	339 (82.9)	24.8		206 (82.7)	27.7		133 (83.1)	20.3	
Premature	70 (17.1)	27.1		43 (17.3)	23.3		27 (16.9)	33.3	
Birth weight			0.367			0.493			0.441
SGA*	50 (12.2)	20		36 (14.5)			14 (8.8)	14.3	
No	359 (87.8)	25.9		213 (85.5)			146 (91.2)	23.3	
Hospitalization			0.475			0.49			0.201
Yes	112 (27.4)	27.7		74 (29.7)	32.4		38 (23.8)	18.4	
No	297 (72.6)	24.2		175 (70.3)	24.6		122 (76.2)	23.8	
Childhood vaccination§			0.339			0.725			0.4
Yes	395 (96.6)	25.6		244 (98.0)	27		151 (94.4)	23.2	
No	14 (3.4)	14.3		5 (2.0)	20		9 (5.6)	11.1	
At least 1 dose of special vaccine&			0.771			0.428			0.569
Yes	51 (12.5)	23.5		33 (13.3)	21.2		18 (11.2)	27.8	
No	358 (87.5)	25.4		216 (86.7)	27.8		142 (88.8)	21.8	
Parent characteristics									
Maternal age			0.276			0.573			0.182
<25	42 (10.3)	35.7		23 (9.2)	39.1		19 (11.9)	31.6	
25-34	267 (65.3)	25.1		162 (65.1)	27.2		105 (65.6)	21.9	
≥35	100 (24.4)	21		64 (25.7)	21.9		36 (22.5)	19.4	
Paternal age			0.987			0.013			0.147
<35	225 (55.0)	28		134 (53.8)	26.9		91 (56.9)	29.7	
≥35	184 (45.0)	21.7		115 (46.2)	27		69 (43.1)	13	
Maternal education			0.002			0.001			0.238
8 years and lessß	202 (49.4)	18.3		135 (54.2)	18.5		67 (41.9)	17.9	
High school and above	219 (50.6)	31.9		114 (45.8)	36.8		93 (58.1)	25.8	
Paternal education			0.024			0.088			0.13
8 years and lessß	186 (45.6)	19.9		115 (46.2)	21.7		71 (44.4)	16.9	
High school and above	223 (54.5)	29.6		134 (53.8)	31.3		89 (55.6)	27	
Health care practices									
Antenatal care visit			<0.001			0.492			<0.001
1-3 times	42 (10.5)	57.1		29 (11.8)	69		13 (8.3)	30.8	
4 and more	359 (89.5)	21.4		216 (88.2)	20.8		143 (91.7)	22.4	
Vaccination during pregnancy			0.012			0.001			0.764
Yes	308 (75.3)	22.1		182 (73.1)	21.4		126 (78.8)	23	
No	101 (24.7)	34.7		67 (26.9)	41.8		34 (21.2)	20.6	

Standard vitamin D supplementation regimen during infancy			<0.001			<0.001			0.377
Yes	371 (90.7)	22.1		227 (91.2)	22.5		144 (90.0)	21.5	
No	38 (9.3)	55.3		22 (8.8)	72.7		16 (10.0)	31.3	
Standard iron supplementation regimen during infancy			0.003			0.004			0.394
Yes	300 (73.3)	21.3		172 (69.1)	21.5		128 (80)	21.1	
No	109 (26.7)	35.8		77 (33.1)	39		32 (20)	28.1	
Standard vitamin D supplementation regimen during pregnancy			<0.001			0.001			0.017
Yes	365 (89.2)	22.2		226 (90.8)	23.9		139 (86.9)	19.4	
No	44 (10.8)	50		23 (9.4)	56.5		21 (13.1)	42.9	
Standard iron supplementation regimen during pregnancy			<0.001			<0.001			0.013
Yes	327 (80.0)	19		201 (80.7)	19.4		126 (78.8)	18.3	
No	82 (20.0)	50		48 (19.3)	58.3		34 (21.2)	38.2	
Vaccine history and ideas of parents									
Childhood vaccination\$ Idea			<0.001			<0.001			<0.001
Protects the child	313 (76.5)	14.4		187 (75.1)	13.4		126 (78.8)	15.9	
I'm hesitant	77 (18.8)	59.7		50 (20.1)	68		27 (16.9)	44.4	
Not safe	12 (2.9)	91.7		9 (3.6)	88.9		3 (1.8)	100	
Refuse	7 (1.7)	14.3		3 (1.2)	0		4 (2.5)	25	
Childhood Vaccination\$ Idea after COVID-19			<0.001			<0.001			<0.001
Yes, necessary	154 (37.7)	12.3		93 (37.4)	11.8		61 (38.1)	13.1	
Yes, hesitant	102 (24.9)	52.9		60 (24.1)	58.3		42 (26.3)	45.2	
No	153 (37.4)	19.6		96 (38.6)	21.9		57 (35.6)	15.8	
Parental COVID-19 vaccination status			<0.001			<0.001			<0.001
Only mother vaccinated	66 (16.1)	19.7		45 (18.1)	22.2		21 (13.1)	14.3	
Only father vaccinated	48 (11.7)	29.2		26 (10.4)	30.8		22 (13.8)	27.3	
Both parents vaccinated	206 (50.4)	14.1		126 (50.6)	15.9		80 (50.0)	11.3	
Neither parent vaccinated	89 (21.8)	52.8		52 (20.9)	55.8		37 (23.1)	48.6	
Maternal COVID-19 vaccination status			<0.001			<0.001			<0.001
Yes	272 (66.5)	15.4		171 (68.7)	17.5		101 (63.1)	11.9	
No	137 (33.5)	44.5		78 (31.3)	47.4		59 (36.9)	40.7	
Paternal COVID19 vaccination status			<0.001			<0.001			0.002
Yes	254 (62.1)	22.5		152 (61.1)	18.4		102 (63.8)	14.7	
No	155 (37.9)	16.9		97 (38.9)	40.2		58 (36.2)	36.2	
Influenced by social media news about vaccines			0.102			0.054			0.009
Yes, hesitant	114 (27.9)	32.5		72 (28.9)	34.7		42 (26.2)	28.6	
Yes, negative	67 (16.4)	32.8		46 (18.5)	30.4		21 (13.1)	38.1	
No, not affect	228 (55.7)	19.3		131 (52.6)	21.4		97 (60.6)	16.5	
Health information sources									
Family elders as a SHIt			0.332			0.213			0.919
Reliable	274 (67.0)	23.7		164 (65.9)	24.4		110 (68.8)	22.7	
Others	135 (33.0)	28.1		85 (34.1)	31.8		50 (31.2)	22	
Pediatrician as a SHIt			0.325			0.35			0.589
Reliable	400 (97.8)	25.5		241 (96.8)	27.4		159 (99.4)	22.6	
Others	9 (2.2)	11.1		8 (3.2)	12.5		1 (0.6)	0	
Family physician as a SHIt			<0.001			<0.001			0.031
Reliable	382 (93.4)	22.8		232 (93.2)	24.1		150 (93.7)	20.7	
Others	27 (6.6)	59.3		17 (6.8)	64.7		10 (6.3)	50	

Family nurse/midwife as a SHI†			0.029			0.109			0.127
Reliable	352 (86.1)	23.3		215 (86.3)	25.1		137 (85.6)	20.4	
Others	57 (13.9)	36.8		34 (13.7)	38.2		23 (14.4)	34.8	
Both members of the family medicine unit as a SHI			<0.001			0.001			0.133
Reliable	344 (84.1)	22.7		208 (83.5)	24.5		136 (85.0)	22.5	
1 of them is reliable	46 (11.2)	28.3		31 (12.5)	25.8		15 (9.4)	33.3	
Others	19 (4.7)	63.2		10 (4.0)	80		9 (5.6)	44.4	
Social media as a SHI†			0.05			0.173			0.145
Reliable	18 (4.4)	5.6		11 (4.4)	9.1		7 (4.4)	0	
Others	391 (95.6)	26.1		238 (95.6)	27.7		153 (95.6)	100	
Government and university website as a SHI†			0.054			0.548			0.029
Reliable	87 (21.3)	17.2		47 (18.9)	23.4		40 (25.0)	10	
Others	322 (78.7)	27.3		202 (81.1)	27.7		120 (75.0)	26.7	
Social media blogger as a SHI†			0.29			0.643			0.222
Reliable	34 (8.3)	17.6		22 (8.8)	22.7		12 (7.5)	8.3	
Others	375 (91.7)	25.9		227 (91.2)	27.3		148 (92.5)	23.6	
Book as a SHI†			0.752			0.283			0.048
Reliable	197 (48.2)	25.9		118 (47.3)	23.7		79 (49.4)	29.1	
Others	212 (51.8)	24.5		131 (52.7)	29.8		81 (50.6)	16	
Television as a SHI†			0.384			0.473			0.685
Reliable	46 (11.2)	30.4		31 (12.4)	32.3		15 (9.4)	26.7	
Others	363 (88.8)	24.5		218 (87.6)	26.1		145 (90.6)	22.1	
TSOY-32 V			0.079						
0-25	84 (43.3)	26.2							
>25-33	58 (29.9)	27.6							
>33	52 (26.8)	11.5							
Disease groups			0.351						
CUA*	249 (60.9)	26.9							
Control	160 (39.1)	22.5							

*=Congenital urogenital anomaly, **= According to the row variable β= Primary and secondary education (8 years), §=Childhood routine vaccination programme †= source of health information, ‡SGA= Small gestational age, §=Childhood routine vaccination programme, ¶= Vaccine Hesitancy Score, α= Upper Q group: ≥25 points in the VHS &=Rota or meningococcal vaccine, †=I'm hesitant, but there's a vaccine tracking system. ‡=Not safe, but there's a vaccine tracking system, †= source of health information, V= Turkey Health Literacy Scale.

DISCUSSION

To the best of our knowledge, this study is the first to evaluate VH specifically among parents of children with CUA.

Our findings reveal that parents in the CUA group who did not routinely administer vitamin D to their infants had 25 times higher odds of VH, suggesting that vaccine refusal may be part of a broader pattern of non-adherence to preventive healthcare. Interestingly, a higher socio-economic and educational profile did not translate to increased vaccine acceptance; mothers with a high school education or higher demonstrated 6.31 times higher odds of VH ($p < 0.001$), and higher TSOY-32 scores were associated with a 4.57-fold increase in hesitancy.

The age of a child also emerged as a significant factor, with parents of children under 24 months showing lower hesitancy rates (15.5%) compared to older age groups ($p = 0.005$), likely due to more frequent healthcare engagement during early infancy. Furthermore, the COVID-19 pandemic significantly reshaped parental attitudes; those who developed hesitancy post-pandemic showed 3.99-fold higher VHS-uQ rates in the CUA group and 11.43-fold higher rates in the CNM group. Conversely, having both parents vaccinated against COVID-19 was strongly associated with lower VH scores in both cohorts ($p < 0.001$), emphasizing that trust in public health interventions remains a cornerstone of routine childhood immunization compliance.

Table 3. Logistic regression analysis of factors affecting vaccine hesitancy in all participants

Features (subgroups (1) vs subgroups)	All			
	Sig.	Exp (B)	95% C.I.	
			Lower	Upper
Disease (CUA vs control)	0.25	1.74	0.68	4.46
Father's education (High school and above vs 8 years and less)	0.001	6.96	2.19	22.14
Child age	0.041			
Child age (24-47 vs <24)	0.087	2.88	0.86	9.67
Child age (≥48 vs <24)	0.015	4.13	1.32	12.97
Father's age (≥35 vs <35)	0.04	0.35	0.13	0.95
Antenatal Care Visit (>4 times vs 1-3 times)				
Standard vitamin D supplementation regimen during pregnancy (No vs Yes)	0.95	1.06	0.17	6.84
Standard iron supplementation regimen during pregnancy (No vs Yes)	0.22	2.56	0.58	11.33
Standard vitamin D supplementation regimen during infancy (No vs Yes)	0.556	1.89	0.23	15.62
Standard iron supplementation regimen during infancy (No vs Yes)	0.401	0.52	0.11	2.42
Childhood Vaccination§ Idea after COVID-19	0.005			
Childhood Vaccination§ Idea after COVID-19 (hesitant vs unaffected)	0.007	5.34	1.58	18.11
Childhood Vaccination§ Idea after COVID-19 (necessary vs unaffected)	0.762	0.83	0.25	2.79
Parental COVID-19 vaccine status	0.002			
COVID-19 vaccine (mother vs none)	0.007	0.11	0.22	0.55
COVID-19 vaccine (father vs none)	0.038	0.18	0.37	0.91
COVID-19 vaccine (both parent vs none)	<0.001	0.08	0.02	0.29
Influenced by social media news about vaccines	0.7			
Influenced by social media news about vaccines (hesitant vs others)	0.498	0.67	0.21	2.16
Influenced by social media news about vaccines (negative vs others)	0.442	0.62	0.18	2.12
Family Nurse/midwife as a SHI† (reliable vs others)	0.658	1.36	0.35	5.37
Government and university website as a SHI† (reliable vs others)	0.382	0.56	0.15	2.06
Book as a SHI† (reliable vs others)				
Vaccination during pregnancy (No vs Yes)	0.123	2.53	0.78	8.21
TSOY	0.056			
(≥33 vs 0-25)	0.034	4.57	1.12	18.65
(25-33 vs 0-25)	0.022	5.52	1.27	23.93
Constant	0.002	0.03		

Health literacy (HL) is a critical determinant in parental vaccine decision-making, as it requires parents to effectively interpret and evaluate complex risk-benefit information regarding their children's health. VH remains a global public health challenge that transcends individual pediatric health, impacting collective societal immunity (22). The literature reveals varying rates of VH worldwide; for instance, a large-scale study involving 4,445 parents reported a 6.1% indecision rate regarding routine vaccinations (23), whereas a study in Nigeria found that 31.6% of participants expressed hesitancy (24). The relationship between HL and vaccination status remains a subject of debate in the international literature. While research in the USA found no significant correlation

between HL and vaccination adherence (25), a similar study in India demonstrated that maternal HL was directly associated with DTP₃ immunization status (26). A notable finding in our study was that VH was 4.57 times higher among participants with higher TSOY-32 scores (≥33), and this hesitancy was significantly more pronounced among those sceptical of childhood vaccination programs ($p < 0.001$). Despite these identified levels of hesitancy, the overall vaccination rate in our cohort remained high at 96.6%. This discrepancy suggests that a robust vaccination tracking system is an effective tool for maintaining high coverage and supporting parental decision-making, even in the presence of underlying hesitancy.

Table 4. Logistic regression analysis of factors affecting vaccine hesitancy in the congenital urogenital anomaly and control group

Features (subgroups (1) vs subgroups)	CUA*				Control			
	Sig.	Exp (B)	95% C.I.		Sig.	Exp (B)	95% C.I.	
			Lower	Upper			Lower	Upper
Maternal education (High school and above vs 8 years and less β)	<0.001	6.31	2.32	17.19				
Paternal education (High school and above vs 8 years and less β)	0.508	1.41	0.51	3.91	0.005	7.52	1.85	30.61
Child age	0.017				0.103			
Child age (24-47 vs <24)	0.396	1.58	0.55	4.56	0.044	3.45	1.03	11.53
Child age ($\geq$ 48 vs <24)	0.005	4.8	1.62	14.19	0.142	2.86	0.7	11.59
Paternal age ($\geq$ 35 vs <35)					0.006	0.21	0.07	0.64
Antenatal care visit (>4 times vs 1-3 times)	0.465	0.6	0.15	2.37				
Standard vitamin D supplementation regimen during pregnancy (No vs Yes)	0.339	0.48	0.1	2.19	0.921	0.91	0.15	5.47
Standard iron supplementation regimen during pregnancy (No vs Yes)	0.409	1.69	0.49	5.89	0.084	4.1	0.83	20.29
Standard vitamin D supplementation regimen during infancy (No vs Yes)	0.001	25.15	4.04	156.5				
Standard iron supplementation regimen during infancy (No vs Yes)	0.938	0.96	0.3	3.05				
Childhood Vaccination§ Idea after COVID-19	<0.001				0.001			
Childhood Vaccination§ Idea after COVID-19 (hesitant vs unaffected)	0.009	3.99	1.42	11.21	0.002	11.43	2.52	51.84
Childhood Vaccination§ Idea after COVID-19 (necessary vs unaffected)	0.005	0.18	0.06	0.6	0.67	0.76	0.21	2.69
Parental COVID-19 vaccine status	<0.001				0.005			
COVID-19 vaccine (mother vs none)	0.002	0.11	0.03	0.45	0.025	0.14	0.02	0.78
COVID-19 vaccine (father vs none)	0.011	0.14	0.03	0.64	0.035	0.19	0.04	0.89
COVID-19 vaccine (both parent vs none)	<0.001	0.07	0.02	0.24	<0.001	0.08	0.02	0.32
Influenced by social media news about vaccines	0.894				0.499			
Influenced by social media news about vaccines (hesitant vs others)	0.639	1.28	0.46	3.52	0.446	0.61	0.17	2.2
Influenced by social media news about vaccines (negative vs others)	0.809	1.16	0.35	3.81	0.249	0.38	0.07	1.98
Family Nurse/midwife as a SHI† (reliable vs others)	0.596	1.41	0.4	4.94				
Government and university website as a SHI† (reliable vs others)					0.27	0.46	0.12	1.82
Book as a SHI† (reliable vs others)					0.907	1.06	0.38	2.97
Vaccination during pregnancy (No vs Yes)	0.384	1.55	0.58	4.15				
Constant	0.195	0.25			0.044	0.17		

*=Congenital Urogenital Anomaly, β = Primary and Secondary Education (8 years), §=Childhood routine vaccination programme, †= source of health information

A woman's journey into motherhood, characterized by a continuous period of learning and adaptation, commences during pregnancy. Throughout this phase, mothers strive to access the most accurate information to make informed and appropriate healthcare decisions for both themselves and their unborn children (27). Enhancing a mother's ability to comprehend health information is critical, as it directly influences her child's long-term well-being and life trajectory. Investing in maternal health literacy and providing access to timely, evidence-based information are essential strategies that yield benefits not only for the individual child but for society as a whole (6,28,29).

Current literature highlights a complex relationship between parental education levels and vaccine confidence. For instance, a study involving 2,214 parents in Rwanda demonstrated that individuals with lower education levels were 3.3 times more likely to express confidence in childhood vaccinations (30). Similarly, research conducted in Saudi Arabia found that VH was 1.71 times higher among those with an education level exceeding high school (31). Our findings align with and reinforce this trend, as we observed that mothers in the CUA group with a high school education or higher had 6.31 times higher odds of VH. This suggests that in more complex clinical contexts like CUA,

higher educational attainment may be associated with increased questioning or heightened sensitivity to vaccine-related information.

The sources of information accessed by parents regarding child health play a pivotal role in shaping their healthcare attitudes and decisions (7,32). In our study, healthcare professionals emerged as the most trusted sources, with parents reporting high levels of reliance on pediatricians (98.0%), family physicians (93.4%), and family medicine midwives (86.7%). This high level of trust in clinical experts is a significant asset for public health, as these professionals serve as the primary line of defence against vaccine misinformation. However, the rapid proliferation of technology, the internet, and social media platforms has created a dual-edged sword: while access to information is easier than ever, distinguishing between evidence-based facts and inaccurate claims has become increasingly challenging for parents, often leading to vaccine hesitancy (32-34). For instance, during the wave of misinformation regarding the MMR vaccine, research indicated that only a quarter of parents consulted their family physicians to verify the information they encountered (35). Similarly, studies on HPV vaccination have shown that relying on information from family and friends can increase the prevalence of insufficient knowledge by 4.34 times (36). Interestingly, our study found that exposure to health information from sources such as social media, family elders, books, television, and public or university websites had no statistically significant effect on vaccine hesitancy in either the CUA or CNM groups, suggesting that for this specific population, clinical trust may outweigh external influences. The COVID-19 pandemic, along with the intense global debates surrounding COVID-19 vaccines, has reignited concerns regarding routine childhood immunization. Factors such as perceived side effects, vaccine ingredients, the nature of vaccine pathogens, institutional distrust, religious or cultural beliefs, and a lack of comprehensive information have historically fueled VH and continue to bring these discussions to the forefront (9,37,38). Conversely, addressing specific concerns of families regarding vaccine safety with evidence-based, scientific answers is recognized as a crucial strategy for restoring and increasing vaccine confidence (39,40). The impact of the pandemic on parental risk perception is further illustrated in studies such as that by Salazar et al., which reported that parents with higher levels of VH tended to be less concerned about the clinical risks of COVID-19 (41). Our findings contribute to this narrative by demonstrating that while VH rates were significantly lower in families where

both parents were vaccinated against COVID-19, exposure to negative vaccine news led to a 3.99-fold increase in the odds of high hesitancy (VHS-uQ) within the CUA group. Based on these results, we suggest that positive outcomes from personal engagement in health practices and the acquisition of reliable health knowledge serve as protective factors, ultimately reducing VH and fostering parental trust in childhood immunization programs.

Strengths and limitations

A primary strength of this study is its multicenter design, encompassing diverse sociocultural regions, which allowed for a more comprehensive evaluation of VH among parents of children with hypospadias and undescended testes. To our knowledge, this is the first study to specifically address VH within the context of CUA. By exploring the unique psychological and clinical landscape of these families, our research provides essential guidance and a novel perspective for physicians and surgeons operating in this specialized field, highlighting the intersection of surgical diagnosis and public health adherence.

Despite these strengths, certain limitations must be acknowledged. First, while multicentric within a national context, the study lacks international scope, which limits the generalizability of the findings to different global socioeconomic and cultural environments. Additionally, the cross-sectional nature of the research precludes a longitudinal assessment of parental attitudes. As the study was conducted post-pandemic, we were unable to utilize a long-term cohort design starting prior to the COVID-19 era, which would have allowed for a more precise temporal analysis of the pandemic's direct impact on shifting vaccination perceptions.

This study identifies significant predictors of VH among mothers of children with CUA. Multivariate analysis reveals that elevated VH is associated with higher maternal education levels and non-adherence to routine infant vitamin D supplementation protocols. Conversely, protective effects were observed in families where both parents were vaccinated against COVID-19 and among those who perceived an increased necessity for childhood vaccinations following the pandemic. These findings suggest that in the specific context of CUA, VH is often intertwined with broader patterns of healthcare engagement and subjective risk assessment.

Clinically, reduced VH correlates strongly with consistent participation in preventive care, the establishment of a robust therapeutic alliance with healthcare providers, and the receipt of evidence-based health education during

clinical encounters. To improve vaccination acceptance in this population, targeted interventions should prioritize three key areas: enhancing provider-parent communication through trusted clinical relationships, delivering science-based immunization information during routine pediatric visits, and developing tailored educational strategies that specifically address the nuanced vaccine concerns of parents with higher health literacy.

Acknowledgements

This study was not supported by any sponsor or funder.

Author Contributions

Conceptualization: S.S.Y and M.S.D.; Methodology: S.S.Y, M.T., and M.S.D.; Software: S.S.Y and M.S.D.; Validation: M.S.D and S.S.Y.; Formal analysis: S.S.Y; Investigation: M.S.D.; Resources: M.T., I.O., A.S.C., D.B., and S.S.; Data curation: M.S.D.; Writing – Original Draft Preparation: M.T., M.S.D, and S.S.Y.; Writing – Review and Editing: M.S.D and S.S.Y.; Visualization: M.S.D. and S.S.Y.; Supervision: S.S.Y.; Project administration: M.T., M.S.D, and S.S.Y. All authors have read and approved the published version of the manuscript.

Statement of Ethics

The study was conducted in accordance with the Declaration of Helsinki and was approved by the Hacettepe University non-interventional Ethics Committee (Approval No. GO 2024/05-42) on January 23, 2024.

Statement of Competing Interest

The authors declare no conflict of interest.

Statement of Data Availability

All data analyzed during this study are included in this published article and available from the corresponding author in the form of anonymized data upon reasonable request.

Statement of Generative AI Use

Not applicable.

Publisher's Note: The statements, opinions, and data contained in AFMN Biomedicine articles are solely those of the individual author(s) and contributor(s) and do not necessarily represent the views of the publisher or the editor(s). The publisher and editor(s) disclaim responsibility for any harm or damage caused by the use of information or products mentioned in the publication.

REFERENCES

1. Bonde JP, Flachs EM, Rimborg S, Glazer CH, Giwerzman A, Ramlau-Hansen CH, et al. The epidemiologic evidence linking prenatal and postnatal exposure to endocrine disrupting chemicals with male reproductive disorders: a systematic review and meta-analysis. *Hum Reprod Update*. 2016;23(1):104-25. [\[CrossRef\]](#)
2. Demirtaş MS, Yalcin SS, Tusat M, Ozmen I, Basar D, Calapoglu AS, et al. Association between air pollution and climate parameters exposure during the periconceptional period and hypospadias risk in Turkish offspring: A case-control study. *Atmospheric Pollution Research* (2025); 102675. [\[CrossRef\]](#)
3. Ben Ari A, Margalit D, Udassin R, Benarroch F. Traumatic Stress among School-Aged Pediatric Surgery Patients and Their Parents. *Eur J Pediatr Surg*. 2019;29(5):437-42. [\[CrossRef\]](#)
4. Gruescu ACS, Popoiu C, Levai MC, Tudor R, Fericean RM, Ravis M. A Cross-Sectional Assessment of Parental Concerns in the Pediatric Surgery Department during the COVID-19 Pandemic. *Healthcare* (Basel). 2023;11(9). [\[CrossRef\]](#)
5. Pawellek M, Kopf FM, Egger N, Dresch C, Mattern U, Brandstetter S. Pathways linking parental health literacy with health behaviours directed at the child: a scoping review. *Health Promot Int*. 2022;37(2). [\[CrossRef\]](#)
6. Banfai-Csonka H, Betlehem J, Deutsch K, Derzsi-Horvath M, Banfai B, Financz J, et al. Health Literacy in Early Childhood: A Systematic Review of Empirical Studies. *Children* (Basel). 2022;9(8). [\[CrossRef\]](#)
7. Cooper S, Schmidt BM, Sambala EZ, Swartz A, Colvin CJ, Leon N, et al. Factors that influence parents' and informal caregivers' views and practices regarding routine childhood vaccination: a qualitative evidence synthesis. *Cochrane Database Syst Rev*. 2021;10(10):CD013265. [\[CrossRef\]](#)
8. Kömürlüoğlu A, Akaydin Gültürk E, Yalçın SS. Turkish Adaptation, Reliability, and Validity Study of the Vaccine Acceptance Instrument. *Vaccines*. 2024;12(5):480. [\[CrossRef\]](#)
9. Morgans HA, Schuster JE, Warady BA. Pediatric Vaccine Hesitancy and COVID-19. *Am J Kidney Dis*. 2023;81(1):13-4. [\[CrossRef\]](#)

10. Demirtaş MS, Artunç NY. Mothers' Perspectives on Childhood Vaccination Programmes after the COVID-19 Pandemic and Subsequent News/Rumours About Vaccination. *Turkish Journal of Pediatric Disease*. 2024;18(6):348-54. [\[CrossRef\]](#)
11. Cataldi JR, O'Leary ST. Parental vaccine hesitancy: scope, causes, and potential responses. *Curr Opin Infect Dis*. 2021;34(5):519-26. [\[CrossRef\]](#)
12. Lafnitzegger A, Gaviria-Agudelo C. Vaccine Hesitancy in Pediatrics. *Adv Pediatr*. 2022;69(1):163-76.. [\[CrossRef\]](#)
13. EuroStat. Statistical Regions in Turkey. 2016.
14. Gonderen Cakmak HS, Uncu D. Relationship between Health Literacy and Medication Adherence of Turkish Cancer Patients Receiving Oral Chemotherapy. *Asia Pac J Oncol Nurs*. 2020;7(4):365-9. [\[CrossRef\]](#)
15. Demirtaş MS, Yaman Artunç N. Mothers' Perspectives on Childhood Vaccination Programmes After the COVID-19 Pandemic and Subsequent News/Rumours About Vaccination. *Turkish J Pediatr Dis* 2024; 18: 348-354. [\[CrossRef\]](#)
16. Bakanlık TCS. Bebek, Çocuk, Ergen İzlem Protokolleri. In: Müdürlüğü HSG, editor. Ankara T.C. Sağlık Bakanlığı Halk Sağlığı Genel Müdürlüğü; 2018.
17. Choi SH, Jo YH, Jo KJ, Park SE. Pediatric and Parents' Attitudes Towards COVID-19 Vaccines and Intention to Vaccinate for Children. *J Korean Med Sci*. 2021;36(31):e227. [\[CrossRef\]](#)
18. Durmaz N, Suman M, Ersoy M, Orun E. Parents' Attitudes toward Childhood Vaccines and COVID-19 Vaccines in a Turkish Pediatric Outpatient Population. *Vaccines (Basel)*. 2022;10(11). [\[CrossRef\]](#)
19. Yalcin SS, Bakacak AG, Topac O. Unvaccinated children as community parasites in National Qualitative Study from Turkey. *BMC Public Health*. 2020;20(1):1087. [\[CrossRef\]](#)
20. Peahl AF, Howell JD. The evolution of prenatal care delivery guidelines in the United States. *Am J Obstet Gynecol*. 2021;224(4):339-47. [\[CrossRef\]](#)
21. Ofliu A, Yalcin SS. Relations of maternal health literacy and parenting practices with early childhood development: a cross-sectional study. *Int J Environ Health Res*. 2024;34(3):1540-50. [\[CrossRef\]](#)
22. Olson O, Berry C, Kumar N. Addressing Parental Vaccine Hesitancy towards Childhood Vaccines in the United States: A Systematic Literature Review of Communication Interventions and Strategies. *Vaccines (Basel)*. 2020;8(4). [\[CrossRef\]](#)
23. Kempe A, Saville AW, Albertin C, Zimet G, Breck A, Helmkamp L, et al. Parental Hesitancy About Routine Childhood and Influenza Vaccinations: A National Survey. *Pediatrics*. 2020;146(1). [\[CrossRef\]](#)
24. Ogbuabor DC, Chime AC. Prevalence and predictors of vaccine hesitancy among expectant mothers in Enugu metropolis, South-east Nigeria. *J Public Health Policy*. 2021;42(2):222-35. [\[CrossRef\]](#)
25. Pati S, Feemster KA, Mohamad Z, Fiks A, Grundmeier R, Cnaan A. Maternal health literacy and late initiation of immunizations among an inner-city birth cohort. *Matern Child Health J*. 2011;15(3):386-94. [\[CrossRef\]](#)
26. Johri M, Subramanian SV, Sylvestre MP, Dudeja S, Chandra D, Kone GK, et al. Association between maternal health literacy and child vaccination in India: a cross-sectional study. *J Epidemiol Community Health*. 2015;69(9):849-57. [\[CrossRef\]](#)
27. Zdolska-Wawrzekiewicz A, Bidzan M, Chrzan-Detkos M, Pizunska D. The Dynamics of Becoming a Mother during Pregnancy and After Childbirth. *Int J Environ Res Public Health*. 2019;17(1). [\[CrossRef\]](#)
28. Ferguson B. Health literacy and health disparities: the role they play in maternal and child health. *Nurs Womens Health*. 2008;12(4):286-98. [\[CrossRef\]](#)
29. Morrison AK, Glick A, Yin HS. Health Literacy: Implications for Child Health. *Pediatr Rev*. 2019;40(6):263-77. [\[CrossRef\]](#)
30. Mbonigaba E, Yu F, Renosa MDC, Cho FN, Chen Q, Denkinger CM, et al. Knowledge and trust of mothers regarding childhood vaccination in Rwanda. *BMC Public Health*. 2024;24(1):1067. [\[CrossRef\]](#)
31. Thirunavukkarasu A, Falji AAM, Al-Hazmi AH, Farhan AB, Alsaidan AA, TA MA, et al. Maternal Perception, Hesitancy, and Satisfaction Toward Childhood Immunization in Primary Health Centers, Hafr Al-Batin: A Multicenter Cross-Sectional Study from Eastern Saudi Arabia. *Risk Manag Healthc Policy*. 2023;16:2357-68. [\[CrossRef\]](#)
32. Demirtas MS, Alici N. The reliability and quality of YouTube videos as a source of breath holding spell. *Ital J Pediatr*. 2024;50(1):8. [\[CrossRef\]](#)
33. Demirtas MS, Alici N. YouTube as a source of information on infantile colic. *Pediatr Int*. 2023;65(1):e15624. [\[CrossRef\]](#)
34. Rosso A, Massimi A, De Vito C, Adamo G, Baccolini V, Marzuillo C, et al. Knowledge and attitudes on pediatric vaccinations and intention to vaccinate in a sample of pregnant women from the City of Rome. *Vaccine*. 2019;37(14):1954-63. [\[CrossRef\]](#)
35. Petts J, Niemeyer S. Health risk communication and amplification: learning from the MMR vaccination controversy. *Health, risk & society*. 2004;6(1):7-23. [\[CrossRef\]](#)
36. Benavidez G, Asare M, Lanning B, Ylitalo K, Fakhoury C, Thompson N, et al. Young adults' human papillomavirus-related knowledge: source of medical information matters. *Public Health*. 2020;182:125-30. [\[CrossRef\]](#)
37. Nuwarda RF, Ramzan I, Weekes L, Kayser V. Vaccine Hesitancy: Contemporary Issues and Historical Background. *Vaccines (Basel)*. 2022;10(10). [\[CrossRef\]](#)
38. Temsah MH, Alhuzaimi AN, Aljamaan F, Bahkali F, Al-Eyadhy A, Alrabiah A, et al. Parental Attitudes and Hesitancy About COVID-19 vs. Routine Childhood Vaccinations: A National Survey. *Front Public Health*. 2021;9:752323. [\[CrossRef\]](#)

39. Arsenault C, Lewis TP, Kapoor NR, Okiro EA, Leslie HH, Armeni P, et al. Health system quality and COVID-19 vaccination: a cross-sectional analysis in 14 countries. *Lancet Glob Health*. 2024;12(1):e156-e65. [\[CrossRef\]](#)
40. Hotez PJ, Batista C, Amor YB, Ergonul O, Figueroa JP, Gilbert S, et al. Global public health security and justice for vaccines and therapeutics in the COVID-19 pandemic. *EClinicalMedicine*. 2021;39:101053. [\[CrossRef\]](#)
41. Salazar TL, Pollard DL, Pina-Thomas DM, Benton MJ. Parental vaccine hesitancy and concerns regarding the COVID-19 virus. *J Pediatr Nurs*. 2022;65:10-5. [\[CrossRef\]](#)